



CS

Dissociative Disorders

Class Objectives

CS

- What are Dissociative Disorders?
 - Dissociative Identity Disorder (DID)
 - Dissociative Amnesia
 - Depersonalization Disorder
 - Derealization Disorder
- Theories and treatments for Dissociative Disorders



Where do you go when you daydream?



Dissociation



☞ Dissociative experiences are somewhere in between a dream state and reality.

☞ Dissociation is a psychological state that produces a lack of connection from the psyche.

- thoughts
- memories
- feelings
- sensations
- identity



Why do we dissociate?



Return to your daydreaming list. When do you zone out or dissociate? How do you feel right before you start daydreaming?

Why do we dissociate?



☞ Even when daydreaming, these dissociative moments result in a temporary *mental escape*.

☞ In situations involving trauma, this provides a break from the fear and pain of the trauma.

- This, in some cases, produces a memory gap surrounding the experience

Depersonalization/ Derealization Disorder



A dissociative disorder in which the individual experiences recurrent and persistent episodes of depersonalization, derealization, or both:

Disconnection from yourself



During episodes of *Depersonalization*, your perception becomes altered so that you temporarily feel they are detached from their own body.

- _____



Disconnection from the world



During an episode of *Derealization* your sense of the

Derealization is a condition in which people feel a sense of unreality or detachment from their surroundings

- _____



I can't remember...



☞ What if we can't remember who we are or how we got to a certain place?? (non-alcohol-related!)

- Amnesia is not just associated with a head trauma, people can have the inability to recall information due to dissociation.

☞ Dissociative amnesia _____

Dissociative Identity Disorder is the same as Schizophrenia.



DID is a VERY controversial psychological disorder.



DID is caused by faulty genetics and biological predispositions



Men are more likely to be diagnosed with DID than women



Dissociative Identity Disorder (DID)



☞ This is one of the most controversial disorders, formerly called Multiple Personality Disorder.

☞ This dissociation disorder involves the creation of "alters", which are partially independent

Separate Identities



☞ People with DID may adopt as many as 100 new identities, all simultaneously coexisting.

- The _____

☞ When under the control of one alter, they may exhibit different behaviors, mannerisms, personalities, gender orientation, and physical properties

- such as handedness, allergies and eyesight

Switch!!



☞ The transition from one alter to another is called a _____

☞ The alter takes control of the host identity and controls their behavior. Due to this, amnesia is required for diagnosis, because sections of time are missing once the alter takes over.

- Some people with DID display "Co-Consciousness" or shared awareness, which varies from person to person.

Who's Effected?



☞ The estimated prevalence of DID in the U.S. population is from 1 in 500 - 1 in 5,000

- Between 250,000 and 2,500,000 people.
- Women are 4 times more likely to be diagnosed
- Research suggests that 3-4% of people hospitalized for psychiatric and drug treatment are affected.

What changes did you see?



☞ When the switch occurs, physical changes are evident, such as

☞ In one study, changes in handedness occurred in 37% of the cases (Putnam et al., 1986).



The person who usually becomes the patient and seeks treatment is the "host" identity.



The alters are usually character-like, often serving specific roles (for example):

- "The Protector"
Handles conflict situations
- "The Whore"
Handles sexuality, sometimes generating income as a prostitute.
- "The child"
Usually represents the age when the child's psyche became fragmented.

It is important to recognize that these



While we see movies and characters with DID portrayed as completely different individuals they are often



Can this disorder be faked?



☞ This is a very controversial disorder and is difficult to determine whether or not the identities are "real" of if the person is faking them for several reasons.

☞ Research supports that persons with DID are

The power of suggestion?



☞ There has been a lot of evidence to suggest that many alters can be created in response to leading questions

- In extreme cases, unethical therapists have encouraged the creation of additional alters by coercion and suggesting false memories.



Can professionals CREATE this disorder??



-Some psychologists believe that MPD is an iatrogenic (physician-induced) behavioral syndrome, promoted by suggestion.

-It is thought by some that DID, like hystero-epilepsy, is created by therapists. This previously rare and disputed diagnosis became popular after the appearance of several best-selling books and movies.

The following is an excerpt taken from a clinician's manual



Review the following slide and think about how this could influence treatment of this disorder

THINK

© Stephen E. Baue, M.D., director of the Dissociative Disorders Treatment Program at a North Carolina hospital.

☞ "It may happen that an alter personality will reveal itself to you during this [assessment] process, but more likely it will not. **So you may have to elicit an alter...** You can begin by indirect questioning such as, 'Have you ever felt like another part of you does things that you can't control?'

☞ "If she gives positive or ambiguous responses ask for specific examples. **You are trying to develop a picture of what the alter personality is like...At this point you may ask the host personality, "Does this set of feelings have a name?"...Often the host personality will not know.** You can then focus upon a particular event or set of behaviors."

Are Clients Just Faking??



☞ Although the high suggestibility of these clients is a factor in the development of alters, research suggests _____

- Miller (1989) confirmed that DID patients display changes in visual acuity, manifest retraction, and eye muscle balance would be difficult to fake.

- Changes in brain function have also been detected by using MRIs to observe brain changes during the time of the switch. Research has shown specific changes in hippocampal and medial temporal areas after the switch (Tsai et. Al, 1999)



The escape into a fantasy world is done to escape the physical and/or emotional pain to survive.

Causes



Research has found that 97-98% of patients had experienced significant trauma, including sexual or physical abuse.

- 68% reported a history of incest

Ross et al., (1990) that 95% of patients reports sexual abuse with a tendency towards extreme, sadistic and often bizarre accounts.

- This research reported documented incidences children being buried alive, tortured with matches, steam irons, razor blades or glass.



Is this like PTSD?



There is a suggested "window of vulnerability" that leads to DID. This theory suggests that trauma prior to age 9 may help explain the development of DID in those with a early history of trauma.

DID can be associated with events such as exposure to combat/war. There is a supported belief that DID is an extreme subtype of PTSD

What Else Contributes DID?

☞ What seems to be most commonly agreed upon in the development of DID is _____

☞ Research has found a high correlation between chaotic, non-supportive family environments and the development of DID after trauma (Waller & Ross, 1997).

Treatment

☞ Symptoms of DID may come and go, but the disorder will not clear up on its own. The process for treatment of DID is not easy or agreed upon by professionals.

☞ The goal of treatment is to integrate the identities into a single identity through long-term psychotherapy, which is usually long and emotionally painful (Ellason et al., 1997).

The prognosis is somewhat unclear. Coon (1986) found that only 5 out of 20 patients achieved full integration of their identities.



Further research showed a 22.2% success for reintegration 2 years after treatment (Ellason, 1997).

New strategies for treatment



Strategies clinicians are using today are based on successful treatments for PTSD, due to commonalities between DID and PTSD (Maladono, 1998).

- The goal for treatment is to identify triggers that provoke memories of trauma/dissociation and neutralize them.
- Most importantly, patients must confront and relive the early trauma so they can gain control (in their mind) over the events (Kuft, 1996).
